SEQUENCE PROTOCOL

| | <110> Roche Diagnostics GmbH | |
|---|--|----------|
| • | <120> Method of identifying N-terminal proBNP | |
| | <130> 51810AWO-SZ | |
| | <140> | |
| | <141> | |
| | <160> 8 | |
| | <170> PatentIn Ver. 2.1 | |
| | <210> 1 <211> 17 | |
| | <212> DNA <213> E. coli | |
| | <400> 1 | |
| | ccggatccca cccgctg | 17 |
| | <210> 2 <211> 79 | |
| | <212> DNA <213> E. coli | |
| | | |
| | <pre><400> 2 cgggatccca cccgctgggt tccccgggtt ccgcttccga cctggaaacc tccggtctgc</pre> | 60 79 |
| | aggaacagcg taaccacct | |
| | <210> 3 <211> 70 | |
| | <212> DNA <213> E. coli | |
| | <400> 3 | |
| | cggttccagg gaggtctgtt caacctgcag ttcggacagt ttaccctgca ggtggttacg | 70 |
| | <210> 4 | |
| | <211> 71 <212> DNA | • |
| | <213> E. coli | |
| | <400> 4 cagacetece tggaaceget geaggaatee eegegteega eeggtgtttg gaaateeegt | - 60 |
| | gaagttgcta c | 7: |
| | <210> 5 <211> 87 | |
| | <212> DNA <213> E. coli | |
| | <400> 5 | |
| | cccaagetta acgeggagea egeagggtgt acagaaceat tttaeggtga ccaeggata | c 6 8 |
| | cttcggtagc aacttcacgg gatttcc | |
| | <210> 6 <211> 19 | |
| | <212> DNA | |

<400> 6 cccaagctta acgcggagc

19

<210> 7 <211> 21 <212> PRT

<213> E. coli

- <400> 7

His Pro Leu Gly Ser Pro Gly Ser Ala Ser Asp Leu Glu Thr Ser Gly

1 5 10 15

Leu Gln Glu Gln Arg 20

<210> 8 <211> 9 <212> PRT <213> E. coli

<400> 8
Glu Leu Gln Val Glu Gln Thr Ser Leu
1 5